

THE POTENTIAL ROLE OF MONOCYTE CHEMOTACTIC AND ACTIVATING
FACTOR (MCAF) IN TUMOR GROWTH

Claus O.C. Zachariae¹, H. Lorraine Thompson¹,
Arthur Anderson², Alberto Mantovani³, Joost J.
Oppenheim¹, and Kouji Matsushima¹.

¹Laboratory of Molecular Immunoregulation, BRMP,
DCT, NCI-FCRF, Frederick, MD 21701-1013, USA,

²USAMRIID, Fort Detrick, Frederick, MD 21701,
USA.

³Istituto di Ricerche Farmacologiche "Mario
Negri", Via Eritrea 62, 20157 Milano, Italy

INTRODUCTION

In the course of the studies of a human monocyte derived neutrophil chemotactic factor (MDNCF/IL 8), a distinct chemotactic factor specific for monocytes (MCF) was also detected. We recently succeeded in purifying MCF from the conditioned media of a myelomonocytic cell line (THP-1) (Matsushima et al., 1989) and cloned the cDNA from a library of an activated human monocytic cell line HL 60 (Furutani et al., 1989). In addition to being chemotactic for monocytes, MCF also activates monocytes to be cytostatic for several types of human tumor cells in vitro. Therefore, we termed this cytokine monocyte chemotactic and activating factor (MCAF).

A variety of murine and human tumor cell lines including our fibrosarcoma cell line 8387 as well as cultured baboon aortic medial smooth muscle cells have been reported to produce MCF-activities (Wang et al., 1986, Benomar et al., 1987, Valente et al., 1988). The in vitro production of this MCF correlated with the macrophage content of in vivo tumors. Furthermore, MCF identical to MCAF has been purified from glioblastoma cell lines and PHA-stimulated PBMC (Yoshimura et al., 1989). In the present paper, we have investigated whether the human fibrosarcoma cell line 8387 produces MCAF. We have also evaluated whether MCAF activates macrophages to produce superoxide anion or lysosomal enzyme release from human monocytes, and established whether MCAF has in vivo monocyte recruitment activity. We have discussed the possibility that these MCAF-induced activities may contribute to tumor resistance.

MATERIALS AND METHODS

Purification of MCAF

Supernatants from fibrosarcoma cell cultures stimulated with 100 ng/ml rTNF α overnight were applied to a heparin Sepharose column. MCAF was eluted with 0.05 M Tris-HCl, 0.5 M NaCl and was chromatographically separated using Sephacryl S-200. The fractions which contained monocyte chemotactic activity were pooled and applied to a carboxymethyl-silica gel column (CM-3SW), which was equilibrated with 0.02 M 3-[N-morpholino] propanesulfonic acid (MOPS), pH 6.5. Finally, samples containing monocyte chemotactic activity were applied to a reverse phase chromatography column (TMS-250 Ultropac column) connected to an HPLC system.

Monocyte Chemotaxis Assay

Monocyte chemotactic activity was measured using a 48-well multiwell Boyden chemotaxis chamber as described (Falk et al., 1980). MCAF was serially diluted in RPMI containing 0.5 % bovine serum albumin (BSA) and added to the lower chamber. Mononuclear cells (2×10^6) free of neutrophils were added to the upper chamber, that was separated from the lower chamber by a polyvinylpyrrolidone-free polycarbonate membrane with a pore size of $8 \mu\text{m}$. After 90 minutes at 37°C the membrane was fixed, dried and stained. The migrated cells consisting of more than 98 % monocytes were counted and the migration was expressed as a chemotactic index (CI), which is the ratio of the number of cells that migrated towards MCAF to the number of cells that migrated towards control medium.

Superoxide Anion Assay

Adherent monocytes were incubated with $10 \mu\text{l}$ of $50 \mu\text{g/ml}$ cytochalasin B in the presence or absence of superoxide dismutase. To each sample, $50 \mu\text{l}$ of the diluted stimuli and $10 \mu\text{l}$ cytochrome C ($120 \mu\text{M}$) was added and the cells incubated for 30 minutes at 37°C . The absorbance of each sample was read in a multiwell ELISA reader at 550 nm. The protein content was measured and the nmoles of superoxide released was calculated using the formula: (absorbance $550 \text{ nm} \times 100$)/($6.3 \times \text{mg cell protein}$).

N-acetyl β -D-glucosaminidase Assay

Adherent monocytes (> 90 % pure) were preincubated with 100 μ l of 5 μ g/ml cytochalasin B, followed by the addition of the appropriate stimulus for 1 hour at 37°C. Supernatants and the lysed cells were added to 400 μ l of the substrate (7.5 mM p-nitrophenyl-acetyl- β -D-glucosamide in 0.125 M citrate buffer containing 0.125 % Triton X-100) for 2.5 hours at 37°C. The amount of enzyme present was defined as that amount of β -N-acetylglucosaminidase that hydrolyzes 1 μ mole of substrate/minute. The percentage of enzyme released was calculated using the formula: enzyme content of supernatant \times 100/(supernatant content + cell monolayer content).

RESULTS AND DISCUSSION

The production of MCAF mRNA upon stimulation of the human fibrosarcoma cell line 8387 with IL 1 and TNF was examined using Northern blotting analysis. Although low expression of MCAF mRNA was found in unstimulated cells, both IL 1 and TNF, but not LPS, increased the level of expression considerably. TNF consistently induced the production of more monocyte chemotactic activity than IL 1. MCAF was, therefore, purified to homogeneity from 8 liters of TNF (100 ng/ml) stimulated fibrosarcoma cells by sequential chromatography including heparin-Sepharose affinity chromatography, CM-HPLC, and RP-HPLC. Material represented by two small absorbance peaks on the CM-HPLC (fraction 25 and 26) that coeluted with monocyte chemotactic activity and β -glucosaminidase releasing activity was applied to RP-HPLC for further purification. In both RP-HPLC eluates a discrete absorbance peak (CM 25, RP 21-22 and CM 26, RP 22-23) coeluted and contained monocyte chemotactic activity, β -glucosaminidase releasing activity and monocyte cytostatic inducing activity suggesting that all these activities are due to a single molecular species. SDS-PAGE analysis of the purified MCAF showed microheterogeneity in relative molecular weight of about 15 kDa. A synthesized fraction of MCAF was used to generate a rabbit polyclonal antibody to MCAF. This antibody reacted in Western blotting analysis with all bands detected on SDS-PAGE. In addition to dose-dependent monocyte chemotactic and N-acetyl β -glucosaminidase releasing activities which are shown in Table 1, purified MCAF also induced superoxide anion release at concentrations above 20 ng/ml.

TABLE 1. Monocyte Chemotactic Activity and N-Acetyl β -Glucosaminidase Releasing Activity of Purified MCAF (CM 25, RP 21-22).

Concentration (pg/ml)	Monocyte Chemotactic Activity (CI)	N-Acetyl β -Glucosaminidase Release (%)
54000	3.91 +/- 0.59	10.0 +/- 0.6
18000	3.47 +/- 0.39	9.1 +/- 0.5
6000	2.87 +/- 0.53	7.3 +/- 0.5
2000	2.68 +/- 0.47	0.4 +/- 0.2
660	1.81 +/- 0.36	-
220	2.00 +/- 0.43	-
73	1.58 +/- 0.24	-
25	1.26 +/- 0.28	-
fMLP (10^{-7} M)	2.54 +/- 0.37	NT
PMA (20 nM)	NT	30.3 +/- 0.1

The identity of the 8387-derived MCAF to MCAF previously purified from the THP-1 cell line was substantiated by specific inhibition of THP-1 derived 125 I-labeled MCAF binding to human PBMC at 37°C by the MCAF enriched unlabeled fractions (CM 25, RP 21-22; CM 26, RP 22-23) as shown in Fig. 1. Both fractions were able to compete with the THP-1 derived MCAF in a dose dependent manner, while another basic polypeptide chemoattractant, IL-8, failed to inhibit the binding of 125 I-labeled MCAF to human PBMC at concentrations from 0.1 ng/ml to 10 μ g/ml. To evaluate the in vivo effects of 8387-derived MCAF, serial dilutions of MCAF in PBS were injected into the ears of Lewis rats at various time points. MCAF showed exclusive monocyte recruitment activity over a dose range from 0.053 ng/ml to 53 ng/ml when assayed at 3, 6 and 18 hours but not at 3 minutes, 30 minutes, or 90 minutes.




Figure 1. Competition for the binding of ¹²⁵I-MCAF to human PBMC by unlabeled purified MCAF. -□- CM25, RP 21-22 (10,6 μg/ml) and -○- CM 26, RP 22-23 (6.8 μg/ml). The binding is expressed as a percentage of the binding obtained with TNP-1 derived ¹²⁵I-MCAF alone.

A large number of tumors are infiltrated with macrophages, which may result in either tumor growth by releasing "growth factors" or exert a tumor suppressive effect through the production of "macrophage activating factors" such as MCAF, IFN-α, TNF, or IL 1. Tumor derived cell lines have been shown to produce MCAF and related molecules (Wang et al., 1986, Benomar et al., 1987, Kuratsu et al., 1989). Such tumors are often infiltrated with macrophages in vivo, and are associated with a better prognosis. IL 1 and TNF can induce MCAF in certain tumor cell types, and in normal fibroblasts and endothelial cells. Thus the production of MCAF by the tumor as well as in neighboring tissues may be regulated by TNF and IL 1. MCAF may contribute to control tumor growth by recruiting activated macrophages to the tumor site. This hypothesis is supported in this paper by the ability of MCAF to induce in vivo monocyte recruitment. Furthermore, MCAF also activates macrophage cytostatic activity and induces enzyme release and O₂⁻ release. The potential utility of these cytokines in cancer treatment remains to be established.

REFERENCES

- Benomar A, Wang JM, Taraboletti G, Ghezzi P, Balotta C, Snyderman R, Dore J-F, Mantovani A (1987). Chemotactic factor and P15E-related chemotaxis inhibitor in human melanoma cell lines with different macrophage content and tumorigenicity in nude mice. *J Immunol* 138:2372-2379.
- Falk W, Goodwin RH, Leonard EJ (1980). A 48-well micro chemotaxis assembly for rapid and accurate measurement of leukocyte migration. *J Immunol Methods* 33:239-247.
- Furutani Y, Nomura H, Notake M, Oyamada Y, Fukui T, Yamada M, Larsen CG, Oppenheim JJ, Matsushima K (1989). Cloning and sequencing of the cDNA for human monocyte chemotactic and activating factor (MCAF). *Biochem Biophys Res Comm* 159:249-255.
- Kuratsu J, Leonard EJ, Yoshimura T (1989). Production and characterization of human glioma cell-derived monocyte chemotactic factor. *JNCI* 81:347-351.
- Matsushima K, Larsen CG, Dubois GC, Oppenheim JJ (1989). Purification and characterization of a novel monocyte chemotactic and activating factor produced by a human myelomonocytic cell line. *J Exp Med* 169:1485-1490.
- Valente AJ, Graves DT, Vialle-Valentin CE, Delgado R, Schwatz C (1988). Purification of a monocyte chemotactic factor secreted by nonhuman primate vascular cells in culture. *Biochem* 27:4162-4168.
- Wang JM, Cianciolo GJ, Snyderman R, Mantovani A (1986). Coexistence of a chemotactic factor and a retroviral P15E-related chemotaxis inhibitor in human tumor cell culture supernatants. *J Immunol* 137:2726-2732.
- Yoshimura T, Robinson EA, Tanaka S, Appella E, Kuratsu J, Leonard EJ (1989). Purification and amino acid analysis of two human glioma-derived monocyte chemoattractants. *J Exp Med* 169:1449-1459.

Molecular and Cellular Biology of Cytokines

*From the Proceedings of the Second International Workshop on
Cytokines. Held at Hilton Head Island, South Carolina,
December 10-14, 1989*

Editors

Joost J. Oppenheim

Laboratory of Molecular Immunoregulation
National Cancer Institute
Frederick Cancer Research Facility
Frederick, Maryland

Michael C. Powanda

Concept Development and
Commercialization Corporation
Mill Valley, California

Matthew J. Kluger

Department of Physiology
University of Michigan Medical School
Ann Arbor, Michigan

Charles A. Dinarello

Department of Medicine
Division of Geographic Medicine and Infectious Diseases
Tufts University School of Medicine
New England Medical Center
Boston, Massachusetts



WILEY-LISS

A JOHN WILEY & SONS, INC., PUBLICATION
NEW YORK • CHICHESTER • BRISBANE • TORONTO • SINGAPORE

Address all Inquiries to the Publisher
Wiley-Liss, Inc., 41 East 11th Street, New York, NY 10003

Copyright © 1990 Wiley-Liss, Inc.

Printed in United States of America

Under the conditions stated below the owner of copyright for this book hereby grants permission to users to make photocopy reproductions of any part or all of its contents for personal or internal organizational use, or for personal or internal use of specific clients. This consent is given on the condition that the copier pay the stated per-copy fee through the Copyright Clearance Center, Incorporated, 27 Congress Street, Salem, MA 01970, as listed in the most current issue of "Permissions to Photocopy" (Publisher's Fee List, distributed by CCC, Inc.), for copying beyond that permitted by sections 107 or 108 of the US Copyright Law. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale.

While the authors, editors, and publisher believe that drug selection and dosage and the specifications and usage of equipment and devices, as set forth in this book, are in accord with current recommendations and practice at the time of publication, they accept no legal responsibility for any errors or omissions, and make no warranty, express or implied, with respect to material contained herein. In view of ongoing research, equipment modifications, changes in governmental regulations and the constant flow of information relating to drug therapy, drug reactions and the use of equipment and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each drug, piece of equipment or device for, among other things, any changes in the instructions or indications of dosage or usage and for added warnings and precautions.

The publication of this volume was facilitated by the authors and editors who submitted the text in a form suitable for direct reproduction without subsequent editing or proofreading by the publisher.

Library of Congress Cataloging-in-Publication Data

International Workshop on Cytokines (2nd : 1989 : Hilton Head Island, S.C.)

Molecular and cellular biology of cytokines : from the proceedings of the Second International Workshop on Cytokines, held at Hilton Head Island, South Carolina, December 10-14, 1989 / editors, Charles A. Dinarello ... [et al.].

p. cm. -- (Progress in leukocyte biology ; v. 10A)

Includes index.

ISBN 0-471-56835-X

I. Cytokines--Congresses. I. Dinarello, Charles A. II. Title. III. Series.

[DNLM: 1. Biological Factors--physiology--congresses. W1 PR6703 v. 10A / QW 568 I618m 1989]

QR185.8.C95I58 1989a

574.87'61--dc20

DNLM/DLC

for Library of Congress

90-12424
CIP